

PREPARING FOR YOUR COLONOSCOPY - SUTAB

Follow the instructions on these papers, **NOT** the booklet that comes in the prep box.

IMPORTANT REMINDERS – 5 DAYS PRIOR

- **5 Days Prior to Procedure:** Please **avoid** high dietary fiber, **seeds, nuts**, corn, popcorn, and iron supplements.
- **Pick Up Supplies 3 Days Prior to Procedure:**
 - ☐ 1 SUTAB bowel prep kit from your pharmacy
 - ☐ Simethicone or Gas-X Tablets
 - ☐ Clear liquids
- **You will need a driver** to take you home after your procedure. You cannot take a bus, taxi, Uber, or Lyft. Your procedure will be cancelled if you do not have a ride.



MEDICATIONS

- **Blood Thinner Medications:**
 - **Stop** Plavix, Aggrenox, Pletal, Prasugrel (Effient) and Ticlid **7 days** prior to procedure.
 - **Stop** Coumadin, Warfarin, or any other blood thinners **5 days** prior to procedure.
 - **Stop** Arixtra, Eliquis, Pradaxa, Xarelto **2-5 days** prior to procedure (depending on renal status).
 - *Please consult with your Cardiologist* about stopping your blood thinning medication and let us know immediately if the time frames listed above are not acceptable.
- **Diabetic Medication**
 - **Insulin twice a day:** inject half your usual dose in the evening before your procedure and take half your usual dose the morning of the procedure.
 - **Insulin only in the morning:** inject your full dose the morning before your procedure but only inject ½ your usual dose the morning of the procedure.
 - Oral Medication: Please hold oral diabetic medication the morning of the procedure.
- **Stop Weight Loss Medication 7 days** prior to procedure which includes GLP1s, Ozempic, Mounjaro, Semaglutide, Ozempic, Wegovey, etc.
- **Continue taking pertinent daily medications** such as heart, seizure, cholesterol, blood pressure, aspirin, etc. with a small sip of water **up until 4 hours** prior to the procedure start time. After this, NOTHING by mouth.

ADDITIONAL NOTES

BOWEL PREP INSTRUCTIONS – SUTAB

1 DAY BEFORE PROCEDURE

Before 9:00AM - You may have a **light breakfast** such as toast, eggs, coffee, tea, non-fat yogurt.

AVOID: meats, cereal, fruits, vegetables, whole grains, oatmeal.

**If you have an afternoon appointment after 1:00pm, you can have a light lunch up until 12:00pm.*


After breakfast, start a **CLEAR LIQUID DIET** up until 4 hours prior to your procedure.

- **Liquids OK to Drink:** Tea, coffee (sweetener only, no cream), water, clear juice, sports drinks (avoid red or purple), broth, ginger ale, sprite, jello (avoid red or purple), popsicles.
- **AVOID:** Anything you can't see through, red/purple liquids, solid food, seeds, alcohol, pulp, nectar drinks, dairy, non/dairy creamers.

Start SUTAB Bowel Prep:

1. **At 6:00pm: Start Dose 1 of SUTAB Bowel Prep**

- a. Fill provided container with 16 oz. of water (to the fill line) and open one SUTAB pill bottle.
- b. Swallow 1 tablet individually every 1-2 minutes with a sip of water. Consume all 12 tablets and 16 ounces of water within 30 minutes.



2. **At 6:30pm: Take 2 Simethicone/Gas-X Tablets**

3. **At 7:30pm: Drink 16 oz of water** over 30 minutes from the container (approximately 1 hour after finishing the first dose)

4. **At 8:30pm: Drink 16 oz of water** over 30 minutes from the container (approximately 30 minutes after finishing last glass of water)

5. **Continue drinking clear liquids** until you go to bed. It is very important to drink plenty of water and other clear liquids throughout the day to avoid dehydration and flush the bowel.



**Nausea, bloating, cramping is normal – pause or slow down the rate of drinking the additional water.
Call the physician office if you have any questions or issues.*

DAY OF PROCEDURE

Take Second Dose of Bowel Prep:

1. **6 hours before your procedure:** start **Dose 2 of SUTAB Bowel Prep**

- a. Fill provided container with 16 oz. of water (to the fill line) and open second SUTAB pill bottle.
- b. Swallow 1 tablet individually every 1-2 minutes with a sip of water. Consume all 12 tablets and 16 ounces of water within 30 minutes.



2. **Take 2 Simethicone/Gas-X Tablets**

3. **You must drink two (2) more 16 oz. containers of water** over the next hour.

4. **You can continue drinking clear liquids** up to 4 hours prior to the procedure



4 Hours before your procedure: **STOP drinking all liquids (including water).**

NOTHING BY MOUTH including gum, candy, or mints. Anything after 4 hours can delay or cancel your procedure.

Check in at the surgery center 1 hour prior to your procedure at .

Bring your ID, insurance card, form of payment, and wear comfortable clothes.