

PLEASE READ ALL ENCLOSED INFORMATION CAREFULLY.

Please complete all enclosed forms and bring them, along with your insurance card(s), copayment or deductible and photo ID on the day of your procedure.

Let us know immediately if your health status or medications change between now and the day of your procedure.

INSURANCE PRE-AUTHORIZATION

If not already done by our office, please check with your insurance company on coverage and to see if pre-authorization is required. **Do not assume this procedure is automatically covered.**

IMPORTANT INSTRUCTIONS REGARDING BLOOD THINNING MEDICATIONS:

7 Days before your procedure: You must stop taking **Plavix, Aggrenox, Pletal and Ticlid**

5 Days before your procedure: You must stop taking **Coumadin, Warfarin, or any other blood thinners**

2-5 days before, depending on renal status, stop **Arixtra, Pradaxa, Xeralto**

Please consult with your Cardiologist about stopping your blood thinning medication ahead of time and let us know immediately if the time frames listed above are not acceptable to your Cardiologist. We will also send a request to your Cardiologist.

IF YOU ARE DIABETIC:

INSULIN: If you inject insulin twice a day, inject ½ your usual dose in the evening before your procedure and take ½ your usual dose the morning of your procedure. If you inject insulin only in the morning, inject your full dose the morning before your procedure but only inject ½ your usual dose the morning of your procedure.

ORAL: Please hold your oral diabetic medication the morning of the procedure.



2 DAYS BEFORE THE PROCEDURE

- Avoid raw fruits, raw vegetables, and grains (especially whole wheat, seeds or nuts)

Colonoscopy: What is a colonoscopy?

Under light sedation, a flexible tube with a light on the end is passed into the rectum and advanced so that the entire colon is examined. It is essential that the colon be very clean so that all polyps or other abnormalities will be seen. If polyps are found, they can usually be removed at the same time. Polyps are important to remove because they can transform into cancer.

Video on Colonoscopy:

[HTTP://WWW.ASGE.ORG/EDUCATION-VIDEOS/COLONVIDEO1.HTML](http://www.asge.org/education-videos/colonvideo1.html)

The MORNING of your procedure

4 hours before your procedure is scheduled
(Actual schedule time, not check-in time)

Remove second bottle of SuFLAVE from the refrigerator and repeat steps 2-3.

MAKE SURE TO COMPLETELY FINISH ALL THE PREP MIXTURE AND WATER TWO HOURS PRIOR TO PROCEDURE TIME.

The day of your procedure

- You are on a clear liquid diet 2 hours before your procedure
- **DO NOT have anything by mouth 2 hours prior to your procedure (this includes water, gum, candy, mints or food items).**



MEDICATIONS THE MORNING OF:

- You may take your blood pressure medications, Parkinson's medications, heart medications and seizure medications with a tiny sip of water.
 - Do not take any other medications for pain, sedatives, anxiety, or injectable blood thinners such as Lovenox. For chronic pain medications, please contact our office at 303-861-0808 for instruction.
 - Diabetic: Insulin – Reduce AM dose by ½ (see Page 1); Oral (ex: Metformin)- Hold the AM of the procedure
- You may brush your teeth. Do not swallow the rinse water.
 - Please don't forget to bring your completed paper work, insurance cards, copayment or any deductible due at time of service.
 - Wear comfortable clothing and shoes. A loose top and sweat pants would be ideal.
 - ***You will need a driver.*** Since you will be receiving sedation, you may not drive until the day after your procedure. This is for your safety. It is the policy of the facility that you have a responsible adult accompany you home after the procedure or your procedure may be cancelled. Call us if you need the number of a bonded transport service.

If you have any questions or concerns about the preparation, please contact us at 303-861-0808.